UNCLASSIFIED

AD NUMBER AD462122 LIMITATION CHANGES TO: Approved for public release; distribution is unlimited. FROM: Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAR 1965. Other requests shall be referred to U.S. Army Natick Laboratories, Natick, MA. AUTHORITY USANL ltr, 23 Sep 1966

UNCLASSIFIED

AD_462122

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION


CAMERON STATION ALEXANDRIA. YIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

162122 CATALOGED BY: DDC

TECHNICAL REPORT ES-15

THE YUMA TEST STATION, ARIZONA THE YUMA TEST STATION, ARIZONA
HOURLY AND DAILY INSOLATION RECORD
1951-1962
DDC

by IVEN BENNETT PH.D. Earth Sciences Division

March 1965

U. S. Army Material Command U. S. ARMY NATICK LABORATORIES Natick, Massachusetts





DOC-IRA

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DDC AVAILABILITY NOTICE

Qualified requesters may obtain copies of this report from Commanding Officer, Defense Documentation Center (DDC) (formerly ASTIA), Cameron Station, Alexandria, Virginia 22314.

DISPOSITION INSTRUCTIONS

Destory; do not return.

AD

TECHNICAL REPORT ES-15

THE YUMA TEST STATION, ARIZONA, HOURLY AND DAILY INSOLATION RECORD 1951 - 1962

by

Iven Bennett, Ph. D.
Desert and Tropic Laboratory
Earth Sciences Division

Project Reference: 1V025001A129

March 1965

U. S. Army Materiel Command
U.S. ARMY NATICK LABORATORIES
Natick, Massachusetts

FOREWORD

Most published studies of the amount of insolation received at the earth's surface are based on energy received over a period of a day. For many purposes, insolation totaled over intervals of time shorter than a day is required. Daily totals compared over the range of latitude found in the United States actually may provide misleading information for some interpretations. For example, in midsummer, due to the greater duration of daylight in northern latitudes and the higher elevation of the sun in southern latitudes, Alaska and Florida will receive essentially the same daily insolation, although instantaneous intensities for much of the day will be considerably higher in Florida. Because of this, data for short periods of time have considerably more meaning for studies of heat load on humans and for some engineering problems.

This report presents descriptive statistics of hourly and daily insolation for Yuma Test Station, Arizona, an environment in which insolation is often extremely high over short time periods. Yuma is also an environment frequently used by the Department of the Army for testing materiel and personnel under conditions of high ambient temperature and high insolation. The data presented here should be of value to environmental testers and engineers concerned with the design of materiel, and to physiologists responsible for defining the limits of human endurance under conditions of high thermal load.

PEVERIL MEIGS, Ph.D. Chief Earth Sciences Division

Approved:

DALE H. SIELING, Ph.D. Scientific Director

W. W. VAUGHAN Brigadier General, USA Commanding

CONTENTS

		Page
List	of Figures	iv
List	of Tables	v
Abst	ract	vi
1.	Introduction	1
2.	Instrumentation and Data	3
	a. The Eppley Pyrheliometer	3
	b. Mountain Standard Times vs. True Solar Time	6
	c. International Pyrheliometric Standard of 1956	6
3.	The Yuma Test Station Climate	6
4.	Presentation of the Data	8
	a. Validity of 1951-1962 Data for Climatic Analysis	8
	b. Hourly Insolation Data	10
	c. Daily Insolation Data	17
5.	Acknowledgments	20
6.	References	23
	ndix: Frequency distribution of hourly insolation for month of the year:	25
	Table II - October 1951 through September 1958	26
	Table III - October 1958 through March 1962	38

LIST OF FIGURES

		Page
1.	Location of Eppley Pyrheliometer at Yuma Test Station, Arizona	2
2.	Close-up of the Eppley Pyrheliometer	2
ځ.	Daily Insolation Traces from an Eppley Pyrheliometer	
	a. Matick, Mass., 2 January 1958 b. Natick, Mass., 4 January 1958	3 3
4.	Comparison of Sunshine and Cloud Cover Monthly Mean Daily Values for 1951-1962 with Values for 1921-1950, U. S. Weather Bureau Airport Station, Yuma, Arizona	9
5.	Histograms of Hourly Insolation, Yuma Test Station, Arisona	
	a. March 1952-1958 b. June 1952-1958 c. September 1952-1958 d. December 1951-1957	12 12 13 13
6.	Mean Hourly Insolation, Yuma Test Station, Arisona, October 1951-September 1958.	14
7.	Mean Hourly Insolation for Half Months, Yuma Test Station, Arizona, October 1951-September 1958.	
	a. Morning Hours b. Afternoon Hours	15 15
8.	Comparison of Distribution of Yuma Hourly Insolation with that of Various Stations.	
	a. June-July b. All Months of Year	16 16
9•	Frequency Distribution by Months of Daily Insolation, Yuma Test Station, Arisona, 20 October 1951-31 March 1962.	21
10.	Daily Values of Insolation, Yuma Test Station, Arisona,	22

LIST OF TABLES

		Page
I	Effect of Ambient Temperature on Performance of Eppley Pyrheliometers (Response is Given in Percent of Response at 80°F)	4
II	Frequency Distributions of Hourly Insolation for Each Month of the Year, Yuma Test Station, Arizona, October 1951 through September 1958 (Hours based on Mountain Standard Time)	
ııı	a. January b. February c. March d. April e. May f. June g. July h. August i. September j. October k. November l. December Frequency Distributions of Hourly Insolation for Each Month of the Year, Yuma Test Station, Arizona,	26 28 99 33 33 35 35 37 35 37
	October 1958 through March 1962 (Hours based on True Solar Time)	
	a. January b. February c. March d. April e. May f. June g. July h. August i. September j. October k. November l. December (1959-1961)	39941244456789 444567899
IV	Yearly Totals and Means of Hourly Insolation in Langleys, Yuma Test Station, Arisona, 1952-1961	17
y	Monthly Totals and Means of Daily Insolation in Langleys, Yuma Test Station, Arisona, 1952-1962	18
VI	Frequency Distribution of Daily Insolation in Langleys for Each Month of the Year, Yumn Test Station, Arizona, 20 October 1951-31 March 1962	19

ABSTRACT

Approximately 10 years of hourly and daily insolation data for the U. S. Army Yuma Test Station, Arizona are summarized in tabular and graphical form. Frequency distributions, totals and means of hourly and daily insolation, are given for each month of the year.

The data were collected by an Eppley pyrhelicmeter (horizontal mount) and an electronic strip chart recorder. Integration and reduction of the data to numerical form were done visually. Information is presented concerning the efficiency of the Eppley pyrheliometer, and of the adequacy of visual data integration and reduction.

As expected from the latitude of the station (32°50'N) and the low incidence of cloud cover, insolation values are high throughout much of the year. From March through October, 30 percent or more of the hours receive at least 60 langleys, and from April through August, 20 percent of the hours receive 80 langleys or more. Intensities of 100 langleys or more occur from April through July, with the maximum during May when 3 percent of the 3 hours centered about True Solar Moon reach 100 langleys or more. Low values occur primarily during early morning, or late afternoon hours. Values less than 20 langleys do not occur at midday except on occasional days in winter and during the summer rainy period of late June and July.

Daily values also run high. Daily totals less than 100 langleys occur only in winter and then infrequently. From April through September, only 1.7 percent of the days have totals less than 400 langleys. On the other hand, 53 percent of the days of these same months receive 700 langleys or more.

THE YUMA TEST STATION, ARIZONA, HOURLY AND DAILY INSOLATION RECORD

1. Introduction

Hourly and daily totals of insolation* have been collected at Yuma. Test Station, Arizona, from 20 October 1951 to the present time. The data have been gathered as part of a comprehensive meteorological support program provided by the Meteorology Department, U.S. Army Electronic Research and Development Activity, to Department of the Army groups testing Army material and personnel at Yuma. In this report a climatic summary is presented of certain features of the record from 20 October 1951 through 31 March 1962. Emphasis is on tabulations of descriptive statistics, and especially on frequency distributions of hourly and daily values of insolation.

Throughout the period of record, the sensor has been an Eppley pyrheliometer (mounted horizontally). Data recording has been by an electronic strip chart recorder.

The following information concerning instrumentation and its location was provided by the Yuma Meteorological Team.

The first data were collected near Building #822 on "Main Post" in a representative area. No instrumentation data are available. Radiation equipment was moved to the "R & D Area" about 1 January 1954. The Eppley pyrheliometer #2297 was then located 430 feet southwest of Building #2700 and 5 feet above the ground. On 19 January 1959, pyrheliometer #3601 was installed on a post 4 feet above the platform on the roof of Building #2700 and 27 feet above the ground (see Fig. 1). Location of the sensor remains the same to date. Exact dates of change are unknown but during 1960 #2293 was used and from 1961 to April 1963 #3554 was used. It is known that #3554 was recalibrated on 3 February 1961.

^{*} In this study, the term "insolation" refers to the total radiant energy from the sun incident on a unit area of a horizontal plane located at the surface of the earth. It is expressed in langleys (1 ly = 1 gm cal/cm/min). It is equivalent to the term "global radiation" (Globalstrahlung) used by many European investigators. It includes all energy in the wavelengths emitted by the sun (approximately 0.1 to 7.0 µ) reaching the bottom of the atmosphere in the direct solar beam and in diffuse sky radiation scattered from the solar beam. It also includes any increment to this total from energy scattered back to the ground by multiple reflection between the ground and the sky or clouds. It does not include energy received at the earth's surface in infrared wavelengths generated by the kinetic energy of molecules of atmospheric gases and particulate matter.




Figure 1. Location of Eppley Pyrheliometer at Yuma Test Station, Arizona




Figure 2. Close-up of Eppley Pyrheliometer

While the pyrheliometer was located on the "Main Post" it was 207 feet above mean sea level. In the "R & D Area" it was first at 325 feet msl, then at 346 feet msl where it remains at present. The approximate latitude and longitude of the instrument during the entire period of record has been 32°50'N and 114°24'W.

2. Instrumentation and Data

a. The Eppley Pyrheliometer

The Eppley pyrheliometer merits a brief description of its general characteristics and of certain sensing and recording difficulties.

Until very recently, no completely reliable system of a non-laboratory nature had been developed for measuring and recording insolation, or for reducing the recorded measurements to digital form*. All insolation data used in the present study were collected by the system used by the United States Weather Bureau, consisting of an Eppley pyrheliometer, an electronic strip-chart recorder, and visual integration of the data. The reliability of this system is typical of most used throughout the world, furnishing, if operated properly, hourly and daily data with an average error of 10 percent or less.

The Eppley pyrheliometer (see Fig. 2) is a thermopile-type instrument that measures insolation by noting the temperature difference created by radiant energy striking a receiving surface consisting of two concentric silver rings, an inner ring covered with lampblack which absorbs virtually all incident insolation, and a white outer ring coated with magnesium oxide with a very high coefficient of reflection for energy in the wavebands of insolation.** Both rings are enclosed in a hermetically sealed spherical glass bulb $3\frac{1}{2}$ inches in diameter. Mounted on the under sides of the rings is the thermopile made up of either 10 or 50 thermocouples. The hot junctions are attached to the black-surfaced disc, the cold junctions are attached to the white-surfaced disc, and the difference in temperature between these junctions generates a small flow of electrons proportional to the intensity of insolation falling on the sensing surfaces.

The spectral response, approximately 0.3 to 5.0μ , of the Eppley to solar energy is due principally to the transmission characteristics of

** The physical description of the instrument given here is taken from Eppley Laboratory, Inc., Bulletin No. 2.

^{*} The Eppley Company, together with the manufacturers of electronic recorders, has recently developed a system that more closely approaches the ideal. This system was not available for use at Yuma and none of the comments in this section apply to it.

the glass used in the bulb. The glass, however, is not controlled for constant transmissivity, and bulbs may vary \$1. htly from one another in the extent of this range. The percent of transmission of the glass varies considerably within the 0.3 to $5.0\,\mu$ range. It is lowest and approaches zero at either end of the range and highest (above 90 percent) between 0.4 and $2.6\,\mu$ where nearly all the energy in insolation is found. On the ultraviolet end the transmission falls rapidly from close to 100 percent at $0.4\,\mu$ to less than 5 percent at $0.28\,\mu$, while through the infrared region transmission remains about 70 percent to about $4.2\,\mu$, after which it declines sharply toward values approaching zero just beyond $5.0\,\mu$.

Laboratory tests have been conducted by several individuals to determine the nature and extent of other errors in the performance of the instrument. One such error involves variations in response with departure of ambient air temperature from the temperature of calibration, 80°F. According to the physical principles governing the electrical output of the thermopile, the output should increase with the ambient temperature. However, the response actually decreases slightly with increasing temperature (MacDonald, 1951, pp. 158-159). Table I, adapted from MacDonald, shows the results of temperature tests with 7 different instruments. The decrease in response with temperature increase above 80°F and conversely, the increase with temperature decrease below 80°F, are apparent. The changes are by no means uniform from instrument to instrument, though they do fall rather closely together. Within the maximum range of temperature normally expected at Yuma, from 40°F to 120°F, the error is less than 4 percent for all the instruments tested.

TABLE I

EFFECT OF AMBIENT TEMPERATURE ON PERFORMANCE OF EPPLEY PYRHELIOMETERS*

(RESPONSE IS GIVEN IN PERCENT OF RESPONSE AT 80°F)

Pyrheliometer**

Ambient Temp. (*F.)	#161 7	#1654	#183 0	#1973	#1977	#1220	#1221	Mean
-40 0 40 80 120	106.2 105.2 102.4 100.0 96.2	102.7 102.5 101.1 100.0 99.0	103.1 102.7 101.1 100.0 97.3	105.4 106.5 104.1 100.0 97.0	105.1 103.0 100.0 96.4	108.1 105.5 102.8 100.0 96.7	103.3 102.4 101.4 100.0 97.8	104.8 104.3 102.3 100.0 97.2

^{*} After a table by MacDonald, 1951 p. 155.

^{**} Data are given for 7 pyrheliometers, rumbered as in column headings.

Another error that must be considered is the so-called cosine error. As the zenith angle of a beam of radiant energy striking the receiving surface of the instrument moves from 0° (normal to the receiving surface) to 90°, assuming constant energy in the beam and no change in other conditions affecting the response of the instrument, the output of the thermopile should decrease with the cosine of the zenith angle, or from 100 percent of the energy in the beam when the sun is directly overhead to O percent when the sun is on the horizon. Laboratory tests by several individuals have shown that this is seldom the case. With low angles of incidence, the instrument generally indicates amounts of insolation that are too small. Miller (1942, p. 324) has suggested as a factor changing coefficients of reflection and absorption, with changing angles of incidence, of the materials used to coat the sensing surface. Woertz and Hand (1941, p. 148) believe the black and white surfaces might not be precisely in the same plane, and that the surfaces might be slightly curved; either of these could cause a variation in response with different angles of incidence. They also discovered that inaccuracies in leveling the surface of the receiver during mounting for field service could contribute to a cosine error of 3.6 percent with zenith angles of 70° or more. Kimball and Hobbs (1923, p. 242) pointed out that the glass envelope could contribute to cosine error unless it was exactly spherical and free from flaws of all kinds.

To give valid data, the Eppley requires constant attention. It is especially important to prevent the accumulation on the outside of the glass bulb of materials such as dust that will cut down transmission of energy to the receiving surface. Dust collection was bound to have been a problem at Yuma, particularly during one period when the instrument was located only 3 or 4 feet above ground level.

A very serious source of possible error involves the electronic recorder used to amplify and record the signal from the pyrheliometer. These recorders are complex and sensitive instruments that require constant expert care to insure proper performance. It is possible for them to drift slowly out of calibration at a rate that would at first escape visual detection. In the present study, many days of data had to be eliminated because values were obviously too high, most likely due to a faulty recorder, and unquestionably much inaccurate data that were not noticeably high or low were left in the study.

A final source of error is produced by the method used to reduce the data to digital form from the line traced by the recorder. Figure 3 contains reproductions of charts similar to those traced at Yuma. In order to calculate the insolation received during any time period on the chart, the area under the curve must be integrated. This is done by eye and involves estimating the mean elevation of the insolation curve for the period in question. Ten-minute periods were used at Yuma, with the values

for these added together to get hourly totals. On an essentially clear day, as shown in Figure 3b, the estimate should be accurate if made with care. On a day with rapidly fluctuating cloud density, Figure 3a, the estimation process becomes very difficult and uncertain. Fortunately, Yuma has a high percentage of clear or nearly clear days; therefore this source of error is held to a minimum.

b. Mountain Standard Time vs. True Solar Time

In the analysis of hourly data for this study, the record has been divided into two periods: (1) 20 October 1951 through 30 September 1958, during which time hourly data were summarized according to Mountain Standard Time; and (2) 1 October 1958 through 31 March 1962, when True Solar Time was the basis for summarization. Hourly data for these two periods are not comparable, since True Solar Noon varies throughout the year in relation to the Standard system. This, of course, does not affect daily insolation values; consequently they have been analyzed for the entire period from 1951 through 1962.

c. International Pyrheliometric Standard of 1956


Effective 1 October 1958, the pyrheliometer at Yuma was recalibrated to conform to the International Pyrheliometric Standard of 1956. Before this, calibration was according to the Smithsonian Standard of 1913. The 1956 standard gives values 2 percent lower than those of the earlier one. Hence in order for all data of the period of record to conform to the 1956 standard, hourly and daily values before 1 October 1958 were reduced by 2 percent.


3. The Yuma Test Station Climate

Yuma Test Station is located in extreme southwestern Arizona in one of the hottest and driest parts of North America. Summers are long and hot, with four months, June through September, having mean daily maximum temperatures above 100°F, and that of July rising to 108°F*. Winters are warm with occasional light frosts. Annual rainfall averages only 3.h inches. The area lies too far south to benefit appreciably from the winter rains of storms originating in the north Pacific Ocean and too far west to gain much from the summer rains associated with moist air from

^{*}All climatic data cited in this discussion are for the Yuma Weather Bureau Airport Station, located 17 miles southwest of the Test Station instrument site. Although climatic data are available for the Test Station, those of the Weather Bureau station are for a considerably longer period of time and are available in summary form. A comparison of some data for the two stations (Nelson, 1957) indicated there is little difference between their climates.

Figure 3





the Gulf of Mexico. September and December receive the most rain, occurring as nighttime thundershowers in September and light continuous rain in December. May and June are the driest months, averaging 0.02 and 0.01 inch respectively.

Most arid areas, though certainly not all, have a low incidence of cloud cover and receive almost the maximum of possible sunshine and insolation. Yuma is not an exception to this. Mean annual cloud cover from sunrise to sunset is only 0.25, and 93 percent of the possible sunshine is received. Much of the cloud cover consists of thin alto-cumulus, altostratus or cirrus clouds that intercept only a small part of incoming solar energy. Maximum cloud cover occurs during the winter months. Monthly mean daily values for December, January, February and March are 0.34, 0.43, 0.32 and 0.34 respectively. June and September have the least cloud cover, 0.12 for each. A slight secondary maximum is found during July and August, with both months averaging 0.23.

The percent of possible sunshine received is high during the entire year. The monthly mean daily value for the least sunny month, January, is 85 percent. All other months, except December with 87 percent, have mean daily values of 90 percent or higher. May, June and September are the sunniest months with mean daily values of 97, 98 and 97 percent. Of the 126 months covered by the present report, 91 percent received 80 percent or more of the possible sunshine and 69 percent received 90 percent or more. Two months, June 1955 and May 1957, received 100 percent of the possible sunshine. Only one month, January 1957 with 64 percent of the possible sunshine, received less than 75 percent.

Days with solid, thick overcast skies that reduce insolation to very low values (100 to 150 ly per day) are rare, whereas completely clear days are common. For the year as a whole Yuma has 6.6 times as many clear as overcast days (Sellers, 1958, p. 23), and many of the overcast days have high, thin clouds that permit passage of considerable solar energy to the earth's surface.


4. Presentation of the Data

a. Validity of 1951-62 Data for Climatic Analysis

The period of record (1951-1962) on which this study is based is relatively short for climatic purposes, hence it would be valuable to determine if the climate during the period of the study is representative of that of a considerably longer period. To make the comparison meaningful to the purpose of this study it is necessary to use climatic elements that correlate highly with insolation. For this purpose, monthly mean daily values of cloud cover and percent of possible sunshine were selected. Figure 4 shows how monthly mean values of these two elements for 1951-1962

COMPARISON OF SUNSHINE AND CLOUD COVER MONTHLY MEAN DAILY VALUES FOR 1951 - 1962 WITH VALUES FOR 1921 - 1950

U.S. WEATHER BUREAU AIRPORT STATION, YUMA, ARIZONA



for the Yuma Weather Bureau Airport Station compare with those of 1921-1950 for the same station. The period 1921-1950 is used by the United States Weather Bureau as a base for establishment of climatic normals. It is apparent that values of the two periods match closely and that insolation statistics for 1951-1962 can be considered reasonably representative of statistics for much longer periods of time.

b. Hourly Insolation Data

The hourly insolation data for Yuma Test Station are summarized in Tables II-IV and Figures 5-8. Through examination of Tables II and III,* the occurrence for any hour in any month of insolation in specified classes of 10 langleys can be determined. Also, by adding percentage frequencies the probability of occurrence above, below, or between certain insolation levels can be estimated.

As expected from the low incidence of cloud cover, hourly insolation values are high throughout much of the year. The 1951-1958 data (Table II) show that from March through October, 30 percent or more of the hours receive at least 60 langleys of insolation. This is a large quantity of radiant energy, equivalent to 1.0 ly per minute, or half the solar constant. From April through August, 20 percent or more of the hourly values are at 80 ly or over. Intensities of 100 ly or more occur from April through July, with the maximum during May when 3 percent of the 3 hours (1200, 1300, 1400)** centered about True Solar Moon reach 100 ly or above. Low values also occur, but primarily during early morning or late afternoon hours when the sum is low in the sky. Low values of 20 ly occur at midday only on occasional days in winter and during the summer monsoon period of late June and July when moist tropical air from the Gulf of Mexico invades the area and causes midday cloudiness.

When all of the hours for a month are considered together, the frequency distribution is generally bimodal (Fig. 5). The two peaks of the distribution are located at or near the extremes of the distribution. This is because there are more hours with low sun and high sun respectively than with intermediate sun, and there is little cloud cover during the day to mask this sun control.

The totals and means of hourly insolation for each year of the period of record are given in Table IV. Means of the 3 high sun hours (1200, 1300, 1400) are 67 ly or higher, even though data from the winter season have been included in their computation. For the 7 hours ending 1000

^{*} Tables of frequency distribution for each month (Tables II and III) are at the end of this report.

^{**} All hourly data are for the hour ending.


TABLE IV


YEARLY TOTALS AND MEANS OF HOURLY INSOLATION IN LANGLETS
YOUR Test Station, Arizona
1952 - 1961

	72	00	00	00	٥ ٦	00	a o	-1 0	00	00	0
	50	240 1	240 1	269 1	354 1	273	267 1	163 1	0.55	೫೦	640
	19	1931 6	2105	2250	2580 8	2334 7	1978 6	179 3 6	913 3	ಕ್ಷ್ಣ	1161 3
	81	5514 18	6483 19	6693 19	80 % 82 %	678k 21	5605 18	5410 17	₹050 11	517 27	4518 14
	11	10696 34	12582 36	12770 37	132TT	12685 39	10443 34	10 20 20 20 20 20 20 20 20 20 20 20 20 20	938k 26	9460 27	₹ 0. %
	91	15872 51	18554 54	18619 53	18698 56	18102 56	15067 49	15695	15405	15437	15704 47
	15	19665 63	22 ₉₁₂	22.732 65	23097	22374 69	18754 61	198 ¹ 3	20210 57	20422 58	2054 68
	#	21772 70	25355 73	253T9 T3	25069 75	24761 TT	21017	22618 71	23811 67	23576 67	23581 71
r Ending	13	22414 72	25669	26196 75	25752	25248 78	21585 71	23566	25288 71	25052 71	24947 T5
Hou	ង	21580	24419 71	24996	24614 74	24007 75	20 699 68	% % %	24749 70	24598	24412 73
	1	1 8842 60	21284	21737 62	21595 65	21027 65	17946 59	20572	22657	2031 8031	1808 1808 1808
	9	14581 47	16480	16616 84	17054 51	163 83 51	14047 46	16 2 33	18328	17954 1.2	17399 47
	8	9333 30	13 80	10767 22	ig w	10TT3	91. 80	34	25.85 35	1240e 35	12491 37
	8	1174 13	13	14 41	58 17	* 3	4157 14	4910 15	6560	6510 81	£°8
	Б	1085	发~	1150	1430	2476	81TT	£~	1896	2008 6	2318
	8	8°	2 °	23 0	Жo	% °	% °	610	153	19	87 T
	YEAR	Total	e e e e	19 19 19 19 19 19 19 19 19 19 19 19 19 1	1975 1946 1970 1970 1970 1970 1970 1970 1970 1970	Total	a see				10 m

NOTE: FROM THE MEDIMINE OF THE RECORD THEOLOGY SEPTEMBER 1958 THE MOURS ARE FOR MOURTAIN STARTED TIME. FROM OCTOBER 1958 TO THE RED OF THE RECORD THEY ARE THUE SOLAR TIME.


HISTOGRAMS OF HCURLY INSOLATION YUMA TEST STATION, ARIZONA





HISTOGRAMS OF HOURLY INSOLATION YUMA TEST STATION, ARIZONA

O = CUMULATIVE PERCENT OF TOTAL HOURS



2000

1900

HOUR ENDING (MOUNTRIN STANDARD TIME)

1200

0000

0800

0700

0000

DEC P.

Š

5

SEP

AUG

₹

₹

MAY

ž

Ž

8

8

8

Figure 6

HOUR ENDING (MOUNTAIN STANDARD TIME)

8

ğ

8

8

1800

1700

1 0041


2000

MEAN HOURLY INSOLATION FOR HALF MONTHS

YUMA TEST STATION, ARIZONA

a. MORNING HOURS
(MOUNTAIN STANDARD TIME)

OCTOBER 1951 - SEPTEMBER 1958



b. AFTERNOON HOURS
(MOUNTAIN STANDARD TIME)

OCTOBER 1951 - SEPTEMBER 1958





Figure 7

COMPARISON OF DISTRIBUTION OF YUMA HOURLY INSOLATION WITH THAT OF VARIOUS STATIONS



through 1600, means are 43 ly or higher. Finally, it can be pointed out that all hours show a uniformity of means from year to year, indicating a high degree of stability of the insolation climate.

Another summary of hourly data is presented in Figure 6, in which the mean insolation of each hour (Mountain Standard Time) by months for the period 1951-1958 are presented. The hour ending 1300 receives the greatest insolation throughout the year. For this hour the mean falls below 50 ly only during December when it is 49.4 ly. The 1300 mean lies above 60 ly from February through October, above 70 ly from March through October, and above 80 ly from April through August, and close to or above 90 ly from May through June. The three hours with highest values average close to 80 ly or more from April through August. Sixty ly is encountered as early in the day as 1000 and as late as 1600 from April to July. The symmetrical arrangement of the isolines about the hour ending 1300 is due to the low incidence of cloud cover and strong sun control at Yuma.

In Figure 7, the annual course of hourly mean insolation is shown for half-month periods. Of interest is the fact that for high sun hours the maximum values occur during the half month 16-31 May, but with the sun lower in the sky during both morning and afternoon the maximum shifts to the first or even the second half of June. For most hours, a dip in values beyond that expected from solar control is indicated for the half month 16-31 July during which time the influx of moist tropical air and associated cloud cover is most pronounced.

Yuma hourly insolation for June-July and the entire year is compared with that of Columbus, Ohio; Sayville, New York; Keyflavik, Iceland; and Wake Island in Figure 8. From Figure 8a it can be seen that during the high sun period (June and July) Yuma values are considerably higher than those of Keyflavik and of the two United States stations, and slightly above those of Wake Island. This is true of Wake Island even though the sun is migher in the sky at Wake during the "high sun' period. From the curves for all months shown in Figure 8b, Yuma again ranks considerably above Keyflavik and the two United States stations and very close to or only slightly below Wake Island. Yuma ranks below Wake Island because of the higher average annual height of the sun at Wake.

c. Daily Insolation Data

The monthly totals and means of daily insolation for every month from January 1952 through December 1962, and the totals and means for each year, and the period of record are presented in Table V.

Table VI contains daily frequency data tabulated by months in class intervals of 50 ly. As with the hourly data, values run high. Daily totals less than 100 ly occur only in winter and then infrequently. Values less than 100 ly represent energy received on heavy overcast days and have

WABLE V

MONTHLY TODALS AND MEANS OF DAILY INSOLATION IN LANGLEYS
Yuma Test Station, Arizona
1952 - 1962

647 680 617 467 377 343 21352 21282 16189 11554 11444 9093 678 686 600 4631 409 303 22031 22267 19124 13772 9964 303 22031 2267 19124 13772 9964 303 19536 2269 1387 6612 13 22031 2267 1662 118 12 22032 226 1662 1436 305 31 22034 12425 1514 413 34 31 22044 15909 16696 1406 340 31 31 2044 15696 1436 1436 340 31 32 2044 15631 17286 14561 340 31 32 2044 16634 14563 14561 340 31 32 2146 16634
2267 19124 13772 9964 654 654 6377 1862 18622 11827 6612 22695 18622 11827 6612 22695 18622 18624 1864 1413 19654
22695 18622 11827 12381 5,22 12381 15,425 19034 16696 14,046 636 756 14,54 18631 17289 14,361 601 776 14,361 18942 14,563 14,361 601 8689 1,207 658 623 4,90 1085 623 1,207 1086 62
25
19094 16696 14-568 9877 636 556 4-548 9877 18631 17285 14-361 10117 601 576 4-63 337 18942 14-563 8112 10311 611 465 262 344 658 623 15-207 10049 658 623 15-207 10049 658 568 4-51 10451 680 568 4-51 10451
18631 17255 14,361 10117 18942 14,563 8112 10311 611 8685 15,207 1004,9 658 623 4,90 335 21088 17037 15211 104,51 680 566 4,91 348
31 30 31 30 611 465 262 344 31 30 31 30 658 623 490 335 21088 17037 15211 10451 680 568 491 348
31 30 31 30 658 623 1,5207 10049 335 335 31 30 31 30 21088 17037 1,5211 10451 680 568 4,91 348
31 30 31 30 21086 17037 15211 10451 680 566 491 348

0.1

			89 899					2 7.0	0.3							93		
			8 5 8 89					15.7	35.5	88.5 2.5	0.3					174		
			750 790 290				18 5.9	135	33.1	81 27.4	12 4.0					342 9.7		
			0 3 t €		_	6.0 9.0	101 38.8	77 26.3	₹8 <u>;</u>	8.75 8.75	99 83	27 27 H				397		
	5		800			21. 6.8	33.8	8,4	11. 9.4	% 17.6	103 34.3	38 13.0	1 0.4			337		
	PREMUENCY DISTRIBUTION OF DAILY INSOLATION IN LANGLESS FOR EACH MONTH OF THE YEAR Young The Station, Artsons 20 October 1951 - 31 March 1962		8 2 5			8,62 29.5	28 10.8	5.1.1	7.1 7.1	19 6.4	57 19.0	8.8 2.5	0.1			28 ε.3		
			8 2 85			88	9.4. 2.	3.0		8 7.7	23 7.7	107 36.6	21.7.7			و ر 7.7		
		삠	8.28	0.3	48 15.7	88.8 18.8	1,5	7.1 7	0.3	9.3	17 5.7	29 9.9	83.5 2.5			2 ⁹⁹ 7.1		
Ľ		in langier	in langer	Class Intervals in Langleys	\$ 3 8	2 0.7	ะหู รั	ส ^ง	2.3		0.3	2.0	4 1.3	2.7	95 35.1	3.0	1 0.3	% 7.2
TABLE VI		SOLATION Nest Stati er 1951 -	term18	8 2 3	30 9.8	23.9	2.6	3.5		2 0.7	5.1	2.3	4 4 H	85.41 14.0	35. 21.6	1.0	239 6.8	
		Class Ir	238	133 33.6	5 .01	3.2	9.0			5.1	3.1.0	2. 0.7	19 7.0	102 33.4	15.3	₹ 8.5		
			5 3 m	73 25.9	17 5.6	1.3				0.3	3.0	9. 7.	1.8	\$. £.3	142 17.2	318 9.0		
			£ 2 €	& %	13 4.3	1.3	1,0				0.3	0.7	5 1.8	35, 11.5	8 17.3	152		
			% 2 % % 2 %	2 6.9	9.E	1.3				0.3	0.3		5.1.8	15,4.9	8 8 6	4°.5		
			28 28	15	5.1.6	0.3			2.0	9.0.	0.3		1 0,4	3.3	9.6	1.3		
			1123	8.6 9.6	0.3	1 0.3						0.3		2 7.0	27.4	25. 8.0		
			४३४	⊬ .	**************************************									0.3	8 7.5	0.5		
			8 3 3	905	50	28	ş	€.	187	\$ 6	8	8	Ę	92	٠	ä		

1. 0.3

8 3 646

¥ \$

ž ž ž

ya,

Jan Jan Apr

virtually no warming effect on the earth's surface or the atmosphere under the cloud layer. At Yuma, only slightly more than 1 day in 100 can be expected to have less than 100 ly total insolation during the winter months (November through February). During summer, low values are seldom seen. For example, from April through September, only 1.7 percent of the days have totals less than 400 ly. On the other hand, 53 percent of the days of these same months receive 700 ly or more. This is truly a massive amount of energy and explains in large part the high daily maximum temperatures, frequently above 100° F, of these months.

Period-of-record monthly means, absolute maxima and minima and certain percentiles (5, 25, 50, 75, 95) of daily insolation are shown in Figure 9. Also presented is the percent of total daily values occurring above the mean. With respect to this percentage, 6 months have 60 percent or more of the days with values larger than the mean, and for all 12 months the figure is in excess of 50 percent. Hence it is not possible to assume, as is so often done, that the mean is a guide to the 50th percentile or median. Also of interest is the considerably smaller frequency range for May than for the other high sun months (June and July). This smaller range is brought about by the complete absence of values below 517 ly during May, whereas the other months experienced at least an occasional day with insolation total less than 200 ly. The frequent occurrence of days with unusually high insolation totals during the high sun season is brought out by the following 75th percentiles for April through August: April, 726 ly; May, 790 ly; June, 814 ly; July, 769 ly; and August, 703 ly.

The annual march of da 'v insolation is presented in Figure 10. Small dots have been placed on the diagram representing the total insolation for each day of the period of record. For comparison, the daily extraterrestrial insolation has been plotted (large dots) for selected days. The distribution of the dots clearly shows the high frequency at all times of the year of values near the maximum possible. Late April and May stand out as a time when low daily totals simply do not occur. Testing under conditions of high thermal load could be done at this time of the year with virtually no danger of interruption due to cloudy weather.

Acknowledgments

Many individuals assisted in the preparation of this report. Mrs. Phyllis Tierney, and Mr. James Hawkins, Computer Branch, respectively, punched the data on cards and supervised the machine analysis. The report was reviewed by Dr. Peveril Meigs, Dr. Llewelyn Williams, and Dr. William Porter. The tables were typed by Miss Carol Christensen. Cartography was performed by Miss Pernel Leuvelink under the supervision of Mr. Aubrey Greenwald.

FREQUENCY DISTRIBUTION BY MONTHS OF DAILY INSOLATION

Yuma Test Station, Arizona, 20 October 1951 - 31 March 1962




Figure 9 21




Figure 10

6. References

Eppley Laboratory, Inc., Newport, Rhode Island, Bulletin No. 2.

Kimball, H. H. and H. E. Hobbs, "A new form of thermoelectric pyrheliometer," Monthly Weather Review, 51(5):239-242. 1923

MacDonald, T. H. "Some characteristics of the Eppley pyrheliometer," Monthly Weather Review, 79(8):153-159. (1951)

Miller, L. F. "Comparative runs between the Eppley and a pyrheliometer having a spherical absorber", Bull Am Meteorol Soc, 23(8):323-328. (1942)

Nelson, R. A., Comparison of Yuma Test Station and Yuma Weather Bureau Meteorological Records, 1952 through 1956, RER 16, Environmental Protection Research Division, Quartermaster Research and Engineering Center, Natick, Mass. November 1957

Sellers, W. D., The Annual and Diurnal Variations of Cloud Amounts and Cloud Types at Six Arizona Cities, Scientific Report No. 8, University of Arizona, Inst Atmospheric Physics. 1958

Woertz, B. B., and I. F. Hand. "The Characteristics of the Eppley Pyrheliometer," Monthly Weather Review, 69 (5):146-148. (1941)

APPENDIX

Frequency Distribution of Hourly Insolation for Each Month of the Year

Table II - October 1951 through September 1958

Table III - October 1958 through March 1962

TABLE II

FREQUENCY DISTRIBUTIONS OF MCURLY INSOLATION FOR EACH MONTH OF YEAR
Yuma Test Station, Arizona
October 1951 through September 1958
(Hours based on Mountain Standard Time)

a. January (1952-1958)

Class Intervals in Langleys

Hour Ending)-9	10-19	20-29	30-39	40-49	50 -5 9	60-69	70-79
06									
07									
08	74	7 ¹ 4 100.0							
09	182	74 40•7	108 59•3						
10	183	13 7.1	29 15.8	92 50-3	49 26. 8				
11	183	3 1.6	19 10.4	15 8.2	43 23.5	91 49.7	12 6.6		
12	183	և 2.Հ	6 3•3	15 8.2	19 10.4	27 14.8	86 47.0	26 14.2	
13	183	2 1.1	8 4.4	13 7.1	13 7.1	26 14.2	66 36.1	51 27. 9	1.2
14	183	0.6	h.h	15 8.2	16 8.7	30 16.4	67 3 6.6	42 23.0	5.5 f
15	183	3 1.6	15 2.2	21 11.5	27 14.8	59. 0	64 35.0	9 4.9	
16	183	12 6.6	23 12.6	35 19.1	66 36.1	46 25.1	0.6		
17	183	23 12.6	76 41.5	79 43.2	5 2.7				
18	172	1 6 0 9 3.0	12 7.0						
19	10	10 100.0							
20									
21									
Totals	1,902	379	304	285	238	264	296	128	8
\$ Freq.		19.9	16.0	15.0	12.5	13.9	15.6	6.7	0.4
Cum. ≸ Freq.		100.0	80.1	64.1	49.1	36.6	22.7	7.1	0.4

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS;
THE LOWER REPRESENTS THE PERCENT OF HOURS.

table II (Cont.)
b. February (1952-1958)
Class Intervals in Langleys

			0100	S THACLAS	10 111 1	P 1 - 1 -				
Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-7 9	80-89
06		•								
07										
08	168	168 100.0								
0 9	185	2 5 1 3. 5	103 55 - 7	57 30.8						
10	185	2 1.1	10 5.4	21	96 51.9	56 30.3				
11	1 8 5		5 2. 7	8 4.3	7 3.8	37 20.0	10 2 55.1	26 14.1		
12	1 8 5	2 1.1	2 1.1	3 1.6	¥ 2.2	13 7.0	13 7.0	111 60.0	3 7 2 0.0	
13	185		3 1.6	1.1	¥ 2.2	9 4.9	10 5.4	5 2 28.1	10 2 55.1	3 1.6
14	185		¥ 2.2	2 1.1	j. 2.2	9 4.9	19 10.3	50 27. 0	93 50.3	\$.2 1
15	185	0.5	3 1.6	6 3.2	3 1.6	10 5.4	41 22 .2	103 55.7	18 9.7	
16	185	1 0.5	7 3.8	7 3.8	2 0 10. 8	65 35.1	79 4 2 .7	6 3. 2		
17	195	5 2.7	12 6.5	52 28.1	103 55•7	13 7.0				
18	185	46 24.9	128 69.2	11 5.9						
19	131	130 99.2	î 0.8							
20										
21										
Totals	2,149	38 0	278	169	241	515	264	348	250	7
\$ Proq.		17.7	12.9	7.9	11.2	9.9	12.3	16.2	11.6	0.3
Cum. \$ Freq.		100.0	82.3	69.4	61.5	50.3	40.4	28.1	11.9	0.3

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE II (Cont.)

c. March (1952-1958)

Class Intervals in Langleys

Hour Frding	No. of Hours	0-9	10-19	20-29	3 0- 39	40.49	50-59	60-69	70-79	80-89	90-99
)6											
07	50	50 100.0									
೧ಕ	185	196 5 7.3	7 4 40.0	5 2.7							
79	187	7 3-7	23 12.3	66 35 • 3	83 44.4	8 4.3					
19	187	0.5	7 3.7	10 5 .3	15 8. 0	62 33.2	81 43.3	10 5• 3	1 0.5		
11	187	0.5	2 1.1	3 1.6	5 2. 7	8 4.3	15 3.0	98 52.4	54 28.9	1	
12	187		3 1.6	1 0.5	3 1.6	4 2.1	11 5.9	11 5. 9	107 57.2	46 24.6	1 0.5
13	187		2 1.1	2 1.1	5 2.7		9 4.8	11 5.9	46 2 4.6	99 5 2 .9	1 3 7.0
14	187		3 1.6	4 2.1	3 1.6	7 3.7	7 3•7	13 7.0	53 28.3	90 48.1	7 3 7
15	1.87		2 1.1	9 4.8	4 2.1	8 4.3	10 5·3	28 15.0	11 2 59.9	14 7.5	
16	187		2 1.1	10 6.4	1 2 8.6	16 27.8	52 49.2	92 1.6	3		
17	187	3 1.6	10 5 • 3	1 8 9.6	46 24.6	103 55.1	7 3.7				
18	187	14 7.5	68 36.4	100 5 3 .5	5 2. 7						
19	180	177 98.3	3 1.7								
2 0											
2 1											
Totals	2,285	359	199	228	181	216	198	263	376	2 50	21
\$ Freq.		15.7	8.7	10.0	7.9	9.5	8.4	11.5	16.5	10.9	0.9
Oum. \$ Freq.		100.0	84.3	75.6	65.6	57 .7	48.2	29.8	28.3	ц.8	0.9

NOTE. OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS:
THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE II (Gont.)
d. April (1952-1958)
Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99	100-109
06	2	2 100.0	-									
σï	257	247 96.1	10 3. 9					•				
ා 8	260	6 2.3	75 28.8	169 6 5. 0	10 3.8							
99	260		6 2.3	14 5.4	45 17.3	179 68. 8	16 6.2					
10	260		2 0.8	4 1.5	12 4.6	8 3.1	43 16.5	184 70.7	7 2.7			
11	26 0		10.4		3 1.2	7 2.7	5 1.9	13 5.0	176 67.7	55 21.2		
12	260			0.4	3 1.2	4 1.5	3 1.2	1.5	7 2.7	199 76.5	39 15.0	
13	26 0				3	2 0.8	3 1.2	5 1.9	o•8 5	74 28.5	169 65.0	2 0.8
14	260			0.4	2 0.8	5 1.9	6 2.3	2 0.8	9 3.5	119 45.8	116 44.6	
15	26 0		3 1.2		4 1.5	3 1.2	6 2.3	6 2.3	49 18.8	189 72.7		
15	26 0		5 1.9	2 0.8	6 2.3	6 2.3	21 8.1	126 48.5	94 36.2			
17	260		5 1.9	9 3•5	19 7.3	105 40.4	122 46. 9					
18	26 0	4 1.5	25 8.8	152 58.5	81 31.2							
19	260	172 66.2	33.8	-								
20	36	36 100.0										
a1 .												
Totals:	3,415	467	218	352	188	319	225	340	344	636	324	P
\$ Freq.		13.7	6.4	10.3	5.5	9.3	6.6	9.9	10.1	18.6	9.5	0.1
Cur:. \$ Freq.		100.0	E6.3	79.9	59.6	64.1	54.8	48.2	3 8.3	28.2	9.6	0,1
W 17 AP 797	M 1980 PT/6	10 TH TH	TR TABLE	9986 11998		THE P	mars or	myme,				

HOTE: OF THE MAINED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUNICIPAL OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE II (Gont.)
d. April (1952-1958)
Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30 -3 9	40-49	50-59	60-69	70-79	80-89	90-99	100-109
06	2	2										
প	257	247 96.1	10 3.9			***				***	- •-	~~
38	26 0	6 2.3	75 28.8	169 65.0)0 3.8							
09	26 0		6 2.3	14 5.4	45 17.3	179 62.8	16 6.2					
10	26 0		2 0.8	4 1.5	12 4.6	8 3.1	43 16.5	184 7 0.7	7 2.7			
11	26 0		0.4		3 1.2	7 2.7	5 1.9	13 5.0	176 67.7	55 21.2		
12	260			0.4	3 1.2	4 1.5	3 1.2	4 1.5	7 2.7	199 76.5	39 15.0	
13	26 0				3	2 0.8	3 1.2	5 1.9	2 0.8	74 28.5	169 65. ວ	o.8
14	26 0			0.4	o.8	5 1.9	6 2.3	o.8	9 3.5	119 45.8	116 44.6	
15	260		3 1.2		4 1.5	3 1.2	6 2.3	6 2.3	49 18.8	189 72.7		
16	26 0		5 1.9	2 0.8	6 2.3	6 2.3	8.1	126 48.5	94 35.2			
17	2 60		5 1.9	9 3.5	19 7.3	105	122 46.9					
18	260	4 1.5	23 8.8	152 58.5	81 31.2							
19	260	172 66.2	38 33.8									
20	36	36 100.0										
21 .												
Totals:	3,415	467	213	352	188	319	225	340	344	636	324	2
\$ Freq.		13.7	6.4	10.3	5.5	9-3	6.6	9.9	10.1	18.6	9.5	0.1
Cum. \$ Freq.		100.0	86.3	79.9	69.6	64.1	54.8	48.2	38.3	26.2	9.6	0.1

HOIS: OF THE MAINED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE II (Cont.)

e. <u>May</u> (1952-1958)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99	100-109
06	75	75 100.0										
٠٠	201	33 40.1	iek 47.7									
08	208	1 0.5	12 5.8	8 3 39•9	112 53.8							
09	208		1 0.5	5 2.4	5 2.4	70 3 3. 6	123 59.1	4 1.9				
10	208			2 1.0	1 0.5	6 2. 9	7 3.4	127 61.1	63 30.3	2 1.0		
11	208						4 1.9	4 1.9	42 30.3	153 73.6	5 2.4	
12	208					1 0.5		3 1.4	7 3.4	89 4 2. 8	105 50.5	3 1.4
13	208			1 0. 5			1 0.5	2 1.0	6 2.9	2 0 9.6	169 81 .2	9 4.3
14	208					2 1.0	1 0.5	2 1.0	9 4.3	35 16.8	153 73.6	6 2 .9
15	208				3 1.4	1.4	1.4	7 3.4	18 8.6	160 76.9	18 8.6	
16	208		1.0 0.5	3		1.9	5 2.4	38 18.3	15 3 73.6	1.9		
17	208		4 1.9	3 1.4	10 4.8	22 10.6	152 73.1	16 7.7		0.5		
13	208		13 6.3	35 16.8	144 69.2	16 7.7						
19 .	208	27 13.0	168 80.8	13 6.3								
20	169	169 100.0										
23	1	1										
Totals	2,948	346	323	145	275	122	294	203	298	464	450	18
\$ Freq.		12.1	11.0	4.9	9.3	4.1	10.0	6.9	10.1	15.7	15.3	0.6
Cum. \$ Proq.		100.0	87.9	76 .9	72.0	62.7	58.	48.6	41.7	31.6	15.9	0.6

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REFRESENTS THE MUMBER OF HOURS; THE LOWER REFRESENTS THE PERCENT OF HOURS.

TABLE II (Cont.)

f. June (1952-1958)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99	100-109
06	123	123 100.0										
প	200	40 20.0	157 78.5	3 1.5								
08	201	1 0.5	9 4.5	48 23.9	141 70.1	2 1.0						
09	201		1 0.5	6 3.0	6 3. 0	58 28.8	127 63.2	3 1.5				
10	201		0.5	¥ 2.0	0.5	2 1.0	0.5	132 65.7	59 29.4	0.5		
11	201		2 1.0	1 0.5	1 0.5		2 1.0	14 2.0	85 42.3	105 52.2	i 0.5	
12	201		2 1.0	1 0.5		0.5	0.5		3 1.5	116 57.7	76 37.8	1 0.5
13	201		2 1.0		1 0.5	1 0.5	1 0.5	2 1.0	3 1.5	60 29. 8	129 64.2	2 1.0
14	201	0.5		1 0.5		2 1.0	1 9.5	1 0.5	14 2.0	66 3 2.8	124 61.7	1 0.5
15	201	0.5	1 0.5			1 0.5	2	0.5	15 7.5	169 84.1	11 5.5	
16	201.	2 1.0			1 0. 5	3 1.5	2 1.0	26 12.9	165 82.1	2 1.0		
17	201.	2 1.0		2.0	3 1.5	11 5.5	122 60.7	58 28.8	1 0.5			
18	501	2 1.0	¥ 2.0	11 5.5	126 62.7	57 28.4	1 0.5					
19	200	9 4.5	111 55.5	80 40.0								
20	196	194 99•0	2 1.0									
21	2	2										
Totals	2,932	377	292	159	2 80	138	26 0	227	335	519	341	4
\$ Freq.		12.9	10.0	5.4	9.6	4.7	8.9	7.7	11,4	17.7	11.6	0.1
Cum. 5 Freq.		100.0	87.1	77.1	71.7	62.1	57.4	48.5	40.8	29.4	11.7	0.1

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS;
THE LOWER REPRESENTS THE PERCENT OF BOURS.

TABLE II (Cont.)

g. <u>July</u> (1952-1958)

Class Intervals in Langleys

Hour Ending	No. of Hours	0- 9	10-19	20-29	30-39	40-45	50-59	60-69	70-79	80-89	90-99	100-109
ત્ ક	44	44 100.0										
17	201	1 41 70.2	59 29.4	1								
n8	505	9	27 13.4	131 64.9	34 16.8	1 0.5						
ù	202	4 2.0	8 4.0	13 6.4	24 11.9	115 56.9	38 18.8					
30	2 02	1 0.5	4 2.0	5 2.5	7 3•5	12 5•9	21 10.4	141 69.8	11 5.4			
11	202	1 0.5		5 2.5	4 2.0	3 1.5	6 3.0	9 4.5	145 71.8	14.4		
12	202	1 0.5	2 1.0	2 1.0	4 2.0	4 2.0	3 1.5	2 1.0	9 4.5	151 74.6	24 11.9	
13	202	2	2 1.0	5 2.5	1 0.5	2 1.0	1 0.5	8 4.0	7 3•5	97 48. o	75 37.1	2 1.0
14	20 2		5 2.5		4 2.0	2 1.0	2	5 2.5	6 3.0	111 55.0	67 33•2	
15	202	2 1.0	¥ 2.0	2 1.0	2 1.0	1 0.5	4 2.0	2 1.0	38 18.8	138 68.3	9 4.5	
1 7	sus	0.5	1.0	3 1.5	1 0. 5	6 3.0	9 4.5	46 22.8	130 64.4	\$.0		
3 *** 1	202		4 2.0	6 3.0	10 5.0	20 9•9	135 66.8	26 12.9	0.5			
13	ละยั	3 1.5	11 5.4	17 8.4	137 67.8	34 16.8						
19	201	25 12•½	132 65.7	44 21.9								
27	194	194 100.0										
71	1	1 100.c										
Totals:	2,863	1,29	260	234	228	200	219	239	347	5 3 0	175	2
% Freq.		15.0	9.1	8.2	8.0	7.0	7.6	8.3	12.3	18.5	6.1	0.1
Cum. 5 Freq.		100.0	85.0	75.9	67.7	59.7	52.7	45.1	36.8	24.7	6.2	0.1

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REFRESENTS THE NUMBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE II (Cont.)
h. August (1952-1958)

Class Intervals in Langleys

Hours Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50 -59	60-69	70-79	80-89	90 -99
06											
07	204	199 97.5	5 2.4								
08	207	11 5.3	70 3 3.8	126 60.9							
99	207	2 1.0	4 2.0	12 5.8	66 31.9	123 59.4					
10	207	2 1.0	2 1.0	5 2.4	8 3.9	15 7·3	81 39·1	94 45.4			
11	207	2 1.0		2 1.0	1 0.5	5 2.4	13 6.3	26 12.6	147 71.0	11 5·3	
12	207		2 1.0	1 0.5		3 1.4	9 4.4	11 5•3	41 19.8	130 62.8	10 4.8
13	207			3 1.4	1 0.5	4 2.0	3 1.4	13 6.3	22 10.7	124 59.9	37 17.9
14	207			2 1.0		6 2.9	6 2.9	11 5·3	20 9.7	140 67.6	22 10 6
15	207		1 C.5	1 0.5	2 1.0	3 1.4	14 6.8	15 7·3	9 5 45.9	75 36.2	1 0.5
16	207	1 0.5	2 1.0	2 1.0	7 3.4	12 5.8	22 10.6	110 53.1	51 2 4.6		
17	207	1 0.5	4 1.9	13 6.3	17 8.2	84 40.6	80 38.6	8 3.9			
18	207	5 2.4	13 6.3	90 43.4	92 44.4	7 3.4					
19	203	76 37.4	120 59.1	7 3.4							
20	130	130 100.0									
21											
Totals:	2,814	429	223	264	194	262	228	288	376	480	70
\$ Freq.		15.2	7.9	9.4	6.9	9.3	8.1	10.2	13.4	17.1	2.5
Cum. \$ Freq.		100.0	84.8	76.9	67 .5	60.6	51.3	43.2	33.0	19.6	2.5

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE HUMBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF BOURS.

TABLE II (Cont.)

1. September (1952-1958)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99
06											
97	187	187 100.0									
08	203	8 3.9	165 81.3	39 14.8							
09	203	0.5	5 2.5	10 4.9	157 77•3	30 14.8					
10	203			0.5	2	26 12.3	15ა 7 6. 8	18 8.9			
11	203					2 1.0	11 5.4	99 48.8	91 44.8		
12	203						3 1.5	19 9.4	110 54.2	71 35.0	
13	203				1 0.5	2 1.0	2	10 4.9	63 31.0	121 59.6	4 2.0
14	203				3 1.5	1 0. 5	5 2.5	18 8.9	87 42.8	89 43.8	
15	203		1 0.5	0.5	3 1.5	1 0.5	12 5.9	66 32.5	115 56.6	5.0 #	
16	203		3 1.5	2 1.0	2 1.0	23 11.3	109 53.7	63 3 1.0	0.5		
17	203	3 1.5	1.0	14 6.9	99 48.8	81 39•9	4 2.0				
18	202	6 3.0	103 51.0	88 43.6	5 2.5						
19	193	185 95.8	8 4.1								
20	9	9 100.0									
21											
Totals	2,621	399	287	146	272	166'	302	293	467	285	Ł.
% Freq.		15.2	10.9	5.6	10.4	6.3	11.5	11.2	17.8	10.9	0.2
Cum. % Freq.		100.0	84. 8	73.9	68.3	57.9	5 1.6	40.1	28.9	11.1	0.2

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

J. October (1951-1957)
Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89
06	1	100.0								
07	46	46 100.0								
08	147	61 41.5	84 57.1	2 1.4						
09	147		6 4.1	73 49•7	66 44.9	2 1.4				
10	147	1 Ø.7	3 2.0	5 3.4	18 12.2	82 55.8	37 25.2	0.7		
11	147		0.7	3 2. 0		16 10.9	30 20.4	91 61.9	6 4.1	
12	147			3 2.0	2 1.4	6 4.1	9 6.1	1 ₁ 1 27.9	83 56.5	3 2.0
13	147			2 1.4	2 1.4	2 1.4	14 9•5	14 9.5	99 67.4	14 9•5
14	147		0.7	2 1.4	4 2.7	6 4.1	14 9•5	42 28.6	7 6 51.7	2 1.4
15	147		2 1.4		3 2.0	19 12.9	42 28.6	75 51.0	6 4.1	
16	147	1 0. 7	4 2.7	6 4.1	30 20.4	85 59.8	20 13.6	0.7		
17	147	3 2.0	25 17.0	8 <u>3</u> 56.5	35 23.8	1 0. 7				
18	144	86 59•7	57 3 9.6	1 0.7						
19	3 6	36 100.0								
20 21										
Totals	1,697	235	183	180	16 0	21 9	166	265	270	19
\$ Freq.		13.9	10.8	10.6	9.4	12.9	9.8	15.6	15.9	1.1
Cum. ≸ Freq.		100.0	86.1	75.3	64.7	55.3	42.4	32.6	17.0	1,1

HOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE PROCEST OF MORE.

TABLE II (Cont.)

k. November (1951-1957)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79
96									
্য	1	100.0							
08	183	177 96 . 7	6 3•3						
09	186	9 4.8	86 46. 2	89 4 7. 8	2 1.1				
10	186	4 2.2	9 4.8	14 7•5	109 58.6	50 26.9			
11	186	1 9.5	5 2.7	8 4.3	10 5.4	?5 40.3	84 45 . 2	3 1.6	
12	186		4 2.2	4 2.2	10 5.4	42 22.6	50 26.9	75 40.3	0.5
13	136		2 1.1	6 3•2	9 4.8	39 21.0	32 17.2	90 48.4	8 4.3
14	186		3 1.6	5 2.7	12 6.4	48 25.8	49 26.3	68 36.6	1 0.5
15	186	1 0.5	6 3 . 2	9 4.8	38 20.4	71 38.2	60 32 . 2	1 0.5	
16	186	2 1.1	15 8.1	119 26 .3	107 57•5	12 6.4	1 0.5		
17	186	21 11.3	133 71.5	31 16.7	1 0.5				
18	174	1 7 3 99.4	0.6						
19									
20									
21									
Totals	2,032	389	270	215	298	337	276	237	10
≸ Freq.		19.1	13.3	10.6	14.7	16.6	13.6	11.6	0.5
Cum. # Freq.		100.0	80.9	67.6	57.0	42.3	25.7	12.1	0.5

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUNBER OF MOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE II (Cont.)

1. December (1951-1957)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79
06									
া									
90	143	143 100.0							
09	191	59 30.9	126 66.0	6 3.1					
10	191	7 3.7	25 13.1	85 44.5	74 38.7				
11	191	4 2.1	6 3. 1	22 11.5	34 17.8	118 61.8	7 3.7		
12	191	2 1.0	9 4.7	9 4 .7	16 8.4	43 22.5	103 53.9	8 4.2	1 0.5
13	191		10 5.2	6 3. 1	18 9.4	37 19.4	96 50.3	20 10.5	4 2.1
14	191	1 0-5	10 5.2	10 5.2	18 9.4	3 7 19.4	106 55.5	6 4.2	1 0.5
15	191	3 1.6	14 7-3	16 8.4	36 18.8	109 57.1	12 6.3	1 10.5	
16	191	9 4. 7	2 3 12.0	69 36.1	82 42.9	7 3•9	1 0.5		
17	191	46 24.1	125 65.4	20 10.5					
18	175	175 100.0							
19	1	100.0							
20									
21									
Totals	2.038	450	348	243	278	351	325	37	6
\$ Freq.		22.1	17.1	11.9	13.6	17.2	16.0	1.8	0.3
Cum. ≸ Freq.		100.0	77.9	60.8	48.9	35-3	18.1	2.1	0.3

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MINGER OF MOURS; THE LOWER REPRESENTS THE PERCENT OF MOURE.

TABLE III

FREQUENCY DISTRIBUTIONS OF HOURLY INSOLATION FOR EACH MONTH OF THE YEAR Yuma Test Station, Arizona October 1958 through March 1962 (Hours Based on True Solar Time)

a. January (1959-1962)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69
06								
07								
08	114	114 100.0						
09	122	20 16.4	98 80.3	¥ 3∙3				
10	122	2 1.6	15 12.3	21 17.2	84 68.8			
11	122	3 2.5	0.8	9.0	10 8.2	91 74.6	6 4.9	
12	122		5 4.1	6 4.9	7 5•7	8 6.6	92 75•4	¥ 3∙3
13	122		7 5•7	0.8	¥ 3∙3	11 9.0	66 54.1	33 2 7. 1
14	122	4 3•3	3 2.5	2 1.6	6 4.9	8.2	88 72.1	9 7.4
15	122	2 1.6	3 2.5	5 4.1	16 13.1	77 63.1	18 14.8	0.8
16	122	i₄ 3∙3	10 8.2	32 26.2	72 59.0	¥ 3•3		
17	122	21 17.2	85 69.7	16 13.1				
18	106	106 100.0						
19								
20								
21								
Totals:	1,318	276	227	98	199	201	270	47
≸ Freq.		20.9	17.2	7.4	15.1	15.3	20.5	3.0
Cum. ≸ Freq.		100.0	79.1	61.9	54.5	39.4	24.1	3.6

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF MURES;
THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

b. February (1959-1962)

Class Intervals in Langleys

Hour Enting	No. of Hours	0-9	10-19	20- 29	30-39	40-49	50-59	60-69	70-79	80-89
06										
07										
08	111	109 98.2	2 1.8							
09	120	7 5.8	61 50.8	52 43.3						
10	120	2 1.7	7 5.8	19 15.8	53 44.2	39 32. 5				
n	120		4 3·3	11 9 .2	6 5.0	21 17.5	65 54.2	13 10.6		
12	120	0.8	2 1.7	5 4.2	7 5.8	9 7•5	19 15.8	63 52.5	14 11.7	
13	120		0.8	8 6.7	5 4.2	6 5.0	13 10.8	48 40.0	38 31.7	0.8
14	120	0.8	2 1.7	6 5.0	3 2.5	10 8.3	10 8.3	60 50.0	28 23.3	
15	120	0.8	3 2.5	4 3∙3	10 8.3	14 11.7	49 40.8	36 30. 0	3 2.5	
16	120	2 1.7	5.2	14 11.7	17 14.2	68 56.7	14 11.7			
17	120	6 5.0	24 20.0	62 51.7	28 23.3					
18	119	81 68. 1	38 31.9							
19	30	30 1 00. 0								
50										
21										
Totals:	1,340	240	149	181	129	167	170	550	83	1
\$ Preq.		17.9	11.1	13.5	9.6	12.5	12.7	16.4	6.2	0.1
Cum. ≸ Freq.		100.0	82.1	71.0	57-5	47.9	35.4	22.7	6.3	0.1

NOTE: OF THE MAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS;

TABLE III (Cont.)

c. March (1959-1962)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60 -6 9	70-79	80-89	90-99
06											
্য	45	45 100.0									
08	90	32 35•6	48 53-3	10 11.1							
09	90	1	3 3•3	31 34.4	43 4 7. 8	12 13.3					
JC.	90	1.1	1.1	3 3-3	7 7.8	24 26.7	45 50.0	9 10 .0			
11	90		1.1	5.5 5	5.5 5	1.1	6 6.7	49 54.4	29 22.2		
12	90				2 .2	3 3•3	1.1	6 6.7	58 64. 4	20.2	
13	90			1.1	1	1.1	5.5 5	8 8.9	27 30.0	49 5 ₹∙4	1
14	90			1.1		# • # #	5.5 5	4 • 4	38 42.2	41 45.6	
15	90		1.1	1.1	3 3•3	#*# #	6 6.7	23 25.6	52 57.8		
16	90	1.1		3 3•3	3 3•3	13 14.4	60 66.7	9 10.0	1.1		
17	90	1.1	5 5•6	15 16.7	59 65.6	10 11.1					
18	90	14 15.6	67 74.4	9 10.0							
19	81	81 100.0									
20											
21											
Totals:	1,116	176	126	76	120	15	122	108	205	110	1
≸ Freq.		15.8	11.3	6.8	10.8	6.4	10.9	9.7	18.4	9.8	0.1
Cum. ≸ Freq.		100.0	84.2	72.9	66.1	55-3	48.9	38.0	28.3	9.9	n •

NOVE: OF THE PAIRED FIGURES IN THIS LABLE, THE UPPER REFRESENTS THE MUNBER OF HOURS; THE LOVE REFRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

d. April (1959-1961)

						_	,,					
				<u>C1</u>	ass Inter	vals in L	angleys					
Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99	100-109
06	22	22 100.0										
07	89	53 59.6	36 40.4									
08	89	1 1.1	10 11.2	42 47.2	36 40.4							
09	89			4 4.5	4 4.5	45 50.6	36 40.4					
10	89	1984 -			2 2.2	4 4.5	5 5.6	54 60.7	24 27.0			
n	89					2 2.2	2 2.2	2 2.2	44 49.4	39 43.8		
12	89				1.1	1 1.1	3.4	2 2.2	4.5	63 70.8	15 16.8	
13	~,							3 3.4	4.5	54 60.7	27 30.3	1.1
14	89					1.1	2 2.2	10 11.2	8 9.0	68 76.4		
15	89				2 2.2	4.5	4 4.5	8 9.0	70 78.6	1 1.1		
16	89	,	•	3 3.4	4 4.5	3 3,4	43 48.3	36 40.4				
17	89	1	3 3.4	3 3.4	44 49.4	37 41.6		1.1				
18	89	2.2 2.2	53 59.6	34 36.2								
19	89	89 100.0										
20												
21												
Totals:	1,179	168	102	86	93	97	95	116	154	225	42	1
\$ Freq.		14.2	8.6	7-3	7.9	8.2	8.1	9.8	13.1	19.1	3.6	0.1
Cum. ≸ Freq.		100.0	85.8	77.2	69.9	62.0	53.8	45.7	35.9	٤٠٩	3.7	0.1

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF BOURS;
THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

e. May (1959-1961)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99	100-109
06	8 3	83 100.0										
ं	85	1 1.2	73 85.9	11 12.9								
08	85		1 1.2	4.7	60 70. 6	20 22.5						
09	8 5			1 1.2	1 1.2	4 4.7	58 68.2	21 24.7				
10	85				1 1.2		1 1.2	16 18.8	65 76.5	2 2.4		
11	85							3 3.5	10 11.8	5 7 67. 1	15 17.6	
12	85							1 1.2	2.4	38 44 .7	¥3 50.6	1.2
13	85						2 2.4	1.2	1.2	23 27.1	50 58.8	8 9.4
14	85						2 2.4	3 3•5	2 2.4	55 64.7	23 27.1	
15	85				1.2	1.2	1.2	2 2.4	49 57.6	31 36.5		
16	85				1.2	2 2.4	12 14.1	69 81.2	1.2			
17	85			4.7	10 11.8	70 82.4	1.2					
18	85	2.4 2.4	12 14.1	71 83.5								
19	84	100.0										
20	1	100.0										
21												
Totals:	1,188	171	86	91	74	97	77	116	130	506	131	9
5 Freq.		14.4	7.2	7.7	6.2	8.2	6.5	9.8	10.9	17.3	11.0	0.8
Cum. ≸ Freq.		100.0	85.6	78.4	70.7	64.5	56.3	49.8	40.0	29.1	u.8	0.8

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUNBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

f. June (1959-1961)

Class Intervals in Langleys

No. of											
Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	90-99	100-109
85	85 100.0										
86	3 3.5	70 8 1.4	13 15.1								
86		1 1.2	2 2.3	69 80.2	14 16.3						
86			1 1.2	1.2	5 5.8	68 79.1	11 12.8				
86					1.2	2 2.3	27 31.4	56 65.1			
86			1 1.2				1 1.2	19 22.1	62 72.1	3 3•5	
86				1 1.2			1 1.2	4.6	49 57.0	31 36.0	4
86				•	1.2	1 1.2	2 2.3		38 44.2	38 44.2	6 7.0
86						2 2.3	3 3•5	1 1.2	52 60.5	28 32.6	
86					1 1.2	2 2.3	1 1.2	47 54.6	35 40.7		
86			1 1.2		2 2.3	5 5.8	64 74.4	14 16.3			
66			2 2.3	6 7.0	45 52.3	33 38.4					
86	3 3.5	3 3•5	50 58.1	30 34.9							
86	48.8 48.8	44 51.2									
51	51 100.6										
1,254	184	118	··• 70	107	69	113	110	141	236	100	6
	14.7	9.4	5.6	8.5	5.5	9.0	8.8	11.2	18.8	8.0	0.5
	100.0	85.3	75.9	70.3	61.8	56.3	47.3	38. 5	27.3	8.5	0.5
	85 86 86 86 86 86 86 86 86 86 86	## 100.6 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.0 ## 100.6	Hours 0-9 10-19 85 85 100.0 86 3 70 3.5 81.4 86 1 1.2 86 86 86 86 86 86 86 86 86 86 81 3.5 3.5 86 42 48 88 51.2 51 51 100.6 1,254 184 118 14.7 9.4	Hours 0-9 10-19 20-29 85 85 100.0 13 86 3.5 81.4 15.1 86 1 2 2.3 86 1 1.2 86 86 1 1.2 86 86 86 86 86 86 86 86 86 86 1.2 86 86 3.5 3.5 58.1 86 48.8 51.2 51 51 51 100.6 10.2 1,254 184 118 70 14.7 9.4 5.6	Hours 0-9 10-19 20-29 30-39 85 85 100.0 13 30-39 86 3 70 13 13 13 13 12 69 12 12 69 12 <	Hours 0-9 10-19 20-29 30-39 40-49 85 85 100.0 13 3.5 81.4 15.1 86 14 15.1 15.1 16.3 16	Hours 0-9 10-19 20-29 30-39 \$60-\$9 50-59 85 85 100.0 13 13.1 14.1 15.1 16.3	Rours 0-9 10-19 20-29 30-39 40-49 50-59 60-69 85 85 100.0 13 15.1 15.1 15.1 15.1 15.1 15.1 15.2 16.3 12.8	Rours 0-9 10-19 20-29 30-39 40-49 50-59 60-69 70-79 85 85 100.0 13 3.5 81.4 15.1 4 <t< th=""><th>Rours 0-9 10-19 20-29 30-39 \$\lambda_{0-49}\$ 50-59 \$60-69 \$70-75 \$80-89 86 3.5 \$81.4 \$15.1 \$13 \$15.1 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 <t< th=""><th> Rours O-9 10-19 20-29 30-39 40-49 50-59 60-69 70-75 80-89 90-99 </th></t<></th></t<>	Rours 0-9 10-19 20-29 30-39 \$\lambda_{0-49}\$ 50-59 \$60-69 \$70-75 \$80-89 86 3.5 \$81.4 \$15.1 \$13 \$15.1 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 \$16.3 \$16.2 <t< th=""><th> Rours O-9 10-19 20-29 30-39 40-49 50-59 60-69 70-75 80-89 90-99 </th></t<>	Rours O-9 10-19 20-29 30-39 40-49 50-59 60-69 70-75 80-89 90-99

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE PENCENT OF MOURS.

TABLE III (Cont.)

g. <u>July</u> (1959-1961)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89	· 90-99	100-109
06	64	64 100.0					•					
07	93	32 34.4	61 65.6									
80	93	5 5.4	9 9.7	35 37.6	44 47•3							
09	93	2 2.2	6 6.4	7 7.5	6.4	34 36. 6	38 40.9					
10	93		1.1	6 6.4	5 5.4	5.5 5	10 10.8	55 59.1	14 15.1			
11	93			4 4.3	1.1	4 4.3	5 5.4	3 3.2	48 51.6	28 3 0.1		
12	93			5.5 5	3 3.2	2.2 2	4 4.3		4 4.3	54 58 . 1	24 25.8	
13	93			1.1	1.1	2.2 2	3 3.2		7 7.5	48 51.6	30 32.3	1 1.1
77	93			1.1	1	1.1	5 5.4	3 3.2	4 4.3	55 59•1	23 24.7	
15	93		1		1.1	1.1	5°5 5	7 7.5	52 55•9	31.2		
16	93			1 1.1	2.2	1.1	8 8.6	68 73.1	12 12.9	1 1.1		
17	93			4 4.3	5 5.4	50 53.8	34 36.6					
18	93	1.1	9 9.7	50 53.8	33 35.5			1				
19	93	36 38.7	57 61.3					***				,
20	65	65 100.0										
21												
Totals:	1,338	205	144	\mathbf{m}_{\cdot}	102	97	109	136	141	215	77	' 1
\$ Freq.		15.3	10.8	8.3	7.6	7.2	8.1	10.2	10.5	16.1	5.8	0.1
Cum. \$ Freq.		100.0	84.7	73.9	65.6	58.0	50.8	42.7	32.5	22.0	5.9	0.1

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS HE NUMBER OF HOURS; THE LOWER REFRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

h. August (1959-1961)

Class Intervals in Langleys

				01000 1.		-11 Tares D.T.					
Hour Ending	No. of Nours	v - 9	10-19	20-29	30-39	40-49	50-5 9	60-69	70-79	80-89	90-99
06	9	9 100.6									
07	93	80 86.0	13 14.0								
08	9 3	6.4	14 15.1	67 72.0	6.4						
09	93	3 3.2	5 5.4	6 6.4	11.8	63 67.7	5 5.4				
10	93		2 2.2	6 6.4	4 4.3	5 5.4	18 19.4	58 62.4			
11	93		2 2.2	1	2 2.2	2 2.2	5 5.4	10 10.8	55 59.1	16 17.2	
12	93			1 1.1		4 4.3	5.4	5 5.4	19 20.4	50 53.8	9 9•7
13	93			1 1.1	1.1	1.1	4 4.3	1.1	11.8	54 58.1	20 21.5
14	93		1.1	1.1		3 3.2	3 3.2	6 6.4	18 19.4	48 51.6	13 14.0
15	93			2 2.2	4 4.3	6 6.4	2 2•2	9 9•7	56 60.2	14 15•1	
16	93	1.1	2 2.2	2 2.2	4 4.3	5 5.4	33 35-5	44 47•3	2 2.2		
17	93	3 3.2	и 4.3	1, 4.3	24 25.8	50 53.8	6.4	1.1	1 1.1		
18	93	10 10.8	26 28.0	55 59.1	2 2.2						
19	92	87 94.6	5 5.4								
20	6	6 100.0									
21											
Totals:	1,223	205	74	146	58.	139	81	134	162	182	42
# Freq.		16.8	6.1	11.9	4.7	11.4	6. 6	u.0	13.2	14.9	3.4
Cum. ≸ Freq.		100.0	83.2	77.1	65.2	60.5	49.1	42.5	31.5	18.3	3.4

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUNDER OF HOURS; THE LOWER REFRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

i. <u>September</u> (1959-1961)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70 -7 9	80-89	90-99
06											
07	88	86 97•7	2 2.3								
08	89		7 7.9	81 91.0	1.1						
09	89		2 2.2	2 2.2	3 3.4	78 87.6	4 4.5				
10	89			1.1	1	3 3.4	18 20.2	65 73.0	1		
11	90		1		1	1.1	1	5 5.6	75 83.3	6 6.7	
12	90	1.1		1.1		1			44 48.9	41 45.6	2 2.2
13	89	1.1		1	1.1	2 2 . 2		1.1	43 48.3	33 37•1	7 7.9
24	89	1 1.1	1		2 2.2	1.1		2.2	60 67.4	21 23.6	1.1
15	89		1.1	2 2.2		2 2.2	1i 12.4	52 58.4	20 22.5	1.1	
16	90		3 3.3	3 3·3	1	47 52.2	3° 33•3	6 6.7			
37	89	2 2.2	8 9.0	44 49.4	28 31.5	6 6.7	1.1				
18	89	45 50.6	42 47.2	1.1	1.1						
19	39	38 9 7. 4	1 2.6								
20	1	100.0									
21											
Totals:	1,111	175	6°	136	39	141	66	131	243	102	10
≸ Freq.		15.8	6.1	12.2	3-5	12.7	7.9	11.8	21.9	9.2	0.9
Cum. \$ Freq.		100.0	84.2	78.1	65.9	62.4	49.7	43.8	32.0	10.1	0.9

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE BUNGER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

J. October (1958-1961)
Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80-89
06										
07	118	118 100.0								
3 8c	124	9 7•3	53 42.7	62 50.0	•					
09	124		8 6.4	4 3.2	56 45 . 2	56 45 . 2				
10	124		0.6	3 2.4	3 2.4	18 14.5	85 68. 6	14 11.3		
11	124		0.8	0.8	3 2.4	3 2.4	11 8.9	76 61.3	29 23.4	
12	124			3 2.4		0.8	7 5.6	42 33.9	. 66 53.2	5 4.0
13	124	0.8		3 2.4	3 2.4	2 1.6	8 6.4	46 37.1	58 46.8	3 2.4
14	124		1 0.8	4 3.2	3 2.4	4 3.2	32 25.8	65 52.4	15 12.1	
15	124	2 1.6	1, 3.2	0.8	9 7.3	55 44.4	47 37•9	6 4.8		
16	124	3 2.4	8 6.4	35 28 . 2	65 52.4	13 10.5				
17	124	30 24.2	81 65.3	13 10.5						
18	85	84 98.8	1 1.2							
19	3	3 100.0							•	
20										
21										
Totals:	1,446	250	158	129	142	152	190	249	168	8
\$ Preq.		17.3	10.9	8.9	9.8	10.5	13.2	17.2	11.6	0.6
Cum. ≸ Freq.		100.0	82.7	71.8	62.9	53.1	42.6	29.4	12.2	0.6

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE FERCENT OF HOURS.

TABLE III (Cont.)

k. November (1958-1961)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59	60-69
06								
07	65	65 100.0						
80	119	37 31.1	80 67 . 2	2 1.7				
09	119	7 5.9	12 10.1	47 39•5	52 43.7	0.8		
10	119	0.8	7 5.9	9 7.6	18 15.1	74 62.2	10 8.4	
11	119		1 0.8	7 5.9	10 8.4	14 11.8	73 61.3	14 11.8
12	119	1 0.8		4 3.4	9 7.6	8 6.7	51 42.9	46 38.7
13	119	1 0.8	1 0.8	6 5.0	8 6.7	9 7.6	60 50.4	34 28.6
14	119	1 0.8	5 4.2	8 6.7	6 5.0	38 31.9	61 51.3	
15	119	0.8	9.2	9 . 2	67 56.3	29 24.4		
16	119	9. [7.6	50 42.0	60 50.4				
17	119	118 99 . 2	1 0.8					
18	3	3 100.0						
19								
20					2			
21					,			
Totals:	1,258	244	168	154	170	173	255	94
\$ Freq.		19.4	13.3	12.2	13.5	13.8	20.3	7.5
Cum. \$ Freq.		100.0	80.6	67.3	55.1	41.6	27.8	7.5

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE MUMBER OF HOURS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

TABLE III (Cont.)

1. December (1958-1961)

Class Intervals in Langleys

Hour Ending	No. of Hours	0-9	10-19	20-29	30-39	40-49	50-59
96							
07	1	100.0					
08	115	113 96.3	2 1.7				
09	119	11 9.2	65 54.6	43 36.1			
10	119	4 3.4	8 6.7	20 16.8	81 68.1	6 5.0	
n	119	2 1.7	7 5 .9	7 5•9	10 8.4	90 75•6	3 2.5
12	77'0	1 0.8	7 5.9	3.4 3.4	10 8.4	17 14.3	80 67.2
13	119		6 5.0	3-4 4	9 7•6	1 ¹ 4 11.8	86 72.3
14	119		7 5•9	7 5•9	15 12.6	71 59•7	19 16.0
15	119	5 4.2	11 9.2	11 9.2	75 63.0	17 14.3	
1.6	119	11 9.2	27 22.7	81 68.1			
17	119	116 97•5	3 2.5				
18	6	6 1.00.0					
19							
20							
21							
Totals:	1,193	270				215	188
\$ Freq.		22.6		14.8		18.0	-
Cum. ≸ Freq.		100.0	77.4	65.4	50.6	33.8	15.8

NOTE: OF THE PAIRED FIGURES IN THIS TABLE, THE UPPER REPRESENTS THE NUMBER OF HAJRS; THE LOWER REPRESENTS THE PERCENT OF HOURS.

Security Classification

U. S. ARMY HATICK LABORATORIES		Unclassified 26 GROUP			
THE YEAR TEST STAFFOR, ARIZONA, HO	URLY AND DAILY INS	MCITALO	RECORD 1951 - 1962		
4. DESCRIPTIVE NOTES (T) report and inclusive dete	•)				
5 AUTHOR(S) (Last nome, first nome, initial)	······································				
BLHHETT, IVEN					
6. REPORT DATE	76. TOTAL NO. OF P	AGES	76. NO. OF REFS		
March 1965	55		7		
88. CONTRACT OR GRANT NO.	96. ORIGINATOR'S REPORT NUMBER(S)				
b. PROJECT NO. 1V025001A129	ES-15				
c.	9b. OTHER REPORT this report)	9b. OTHER REPORT NO(5) (Any other numbers that may be assigned this report)			
d.					
11. SUPPLEMENTARY NOTES	Laboratory, Ea	rth Sci	vity Desert and Tropic ences Division, U. S. eies, Hatick, Mass.		
Approximately 10 years of hourly a Test Station, Arizona, are summarize distributions, totals and means of hourth of the year.	ed in tabular and g nourly and daily in	raphica solatio	d form. Frequency on, are given for each		
The data were collected by an Lopl electronic strip chart recorder. In form were done visually. Information Eppley pyrheliometer, and of the address expected from the latitude of the strip of the stri	ntegration and redu on is presented cor equacy of visual de	ection on ecerning eta inte	of the data to numerical the efficiency of the egration and reduction.		
As expected from the latitude of the cloud cover, insolation values are he through October, 30 percent or more from April through August, 20 percent Intensities of 100 langleys or more during May when 3 percent of the 3 he langleys or more. Low values occur	nigh throughout mud of the hours receint of the hours red occur from April the nours centered about	th of the live at 1 the live 80 through at True	ne year. From March least 60 langleys, and langleys or more. July, with the maximum Solar Noon reach 100		

DOCUMENT CONTROL DA : - - R&D
(Security classification of titus, body of abstract and indexing annotation must be entered when the overall report is classified)

Daily values also run high. Daily totals less than 100 langleys occur only in winter and then infrequently. From April through September, only 1.7 percent of the days have totals less than 400 langleys. On the other hand, 53 percent of the days of these same months receive 700 langleys or more.

noon hours. Values less than 20 langleys do not occur at midday except on occasional days in winter and during the summer rainy period of late June and July.

DD 1508M 1473

Unclassified
Security Classification

Security Classification

KEY WORDS	LINK A		LINK B		LINK C	
KET HORUS	ROLE	WT	ROLE	wt	ROLE	WT
Solar radiation	8					
Insolation	8					
Climatology	1 4]			
Radiation measurement systems	10					
Desert tests	4	ŀ	1		1	
Pyrheliometers	10					
Data	8				}	
Electronic recording systems	10					
Yuma, Arizona	9					
Deserts	4	1			1	
]		1		1	
					1	i

INSTRUCTIONS

- 1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of L \sim fense activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases ahould be uncleasified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a apecific reporting period is covered.
- 5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of aervice. The name of the principal author is an absolute minimum requirement.
- 6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- 8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, auch as project number, subproject number, system numbera, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the snonsor), also enter this number(s).

- 10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further disaemination of the report, other than those imposed by security classification, using atandard atatements such as:
 - "Qualified requesters may obtain copies of this report from DDC."
 - (2) "Foreign announcement and diasemination of thia report by DDC is not authorized."
 - (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
 - (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
 - (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, Indicate this fact and enter the price, if known.

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory aponaoring (paying for) the reaearch and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a hrief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports he unclassified Each paragraph of the abstract shall end with an indicati of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abatract. However, the suggested length ia from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or ahort phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Idenfiers, such as equipment model deaignation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

Unclassified

Security Classification